



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

Remarks on Problem 179 by TSURUICHI HAYASHI, Koto Shiha Gakko, Tokyo, Japan.

The following question proposed by Dr. L. E. Dickson, The University of Chicago, remains unsolved in Vol. XII, 1905, p. 238:

Find the roots of the algebraically solvable quintic equation

$$x^5 + qx^2 + px + \frac{1}{5} \left[\frac{q^2}{p} - \frac{p^3}{5q} \right] = 0.$$

I think that the coefficients of x^2 and x must be interchanged and thus the equation must become

$$x^5 + px^2 + qx + \frac{1}{5} \left[\frac{q^2}{p} - \frac{p^3}{5q} \right] = 0.$$

If so, the roots are represented by

$$-\omega_{\lambda^5} \sqrt[5]{\frac{q^2}{5p}} + \omega_{\lambda^3} \sqrt[5]{\frac{p^3}{25q}}$$

where $\lambda=1, 2, 3, 4, 5$, and $(\omega_{\lambda})^5=1$.

Remark by the PROPOSER.

The problem was printed incorrectly; it should have read $x^5 + px^2 + qx + \dots$, with the letters p, q in their natural order. Since p is of the third degree in the roots, and q of the fourth, the constant term is of the fifth degree, as should be the case.

Mr. T. Hayashi's solution of the corrected equation has, doubtless by an oversight in copying, the terms ω and ω^3 interchanged.

Solution. Since the terms x^4 and x^3 are lacking, the simplest expressions to assume for the roots are

$$r_{\lambda} = \omega_{\lambda^3} A + \omega_{\lambda} B \quad (\lambda=1, \dots, 5; \omega_{\lambda^5}=1).$$

Then by $\sum_{\lambda=1}^5 \omega_{\lambda^r} = 0$ for $r=1, \dots, 4$, we have for $S_t = \sum r_{\lambda}^t$,

$$S_1 = S_2 = 0, \quad S_3 = 15AB^2, \quad S_4 = 20A^3B, \quad S_5 = 5A^5 + 5B^5.$$

But for $x^5 + px^2 + qx + r = 0$,

$$S_1=S_2=0, \quad S_3=-3p, \quad S_4=-4q, \quad S_5=-5r.$$

$$\text{Hence, } A = -\sqrt[5]{\frac{q^2}{5p}}, \quad B = \sqrt[5]{\frac{p^3}{25q}}, \quad r = \frac{q^2}{5p} - \frac{p^3}{25q}.$$

276. Proposed by W. J. GREENSTREET, M. A., Editor of *The Mathematical Gazette*, Stroud, England.

If x_1, x_2, \dots, x_n be unequal, and $f(x)$ be a rational integral function of degree $\geq n-2$, then shall

$$\sum_{r=1}^{r=n-1} \frac{f(x_r)}{(x_r-x_1)(x_r-x_2)\dots(x_r-x_n)} = 0.$$

Solution by the PROPOSER.

The left hand side written at length is

$$\begin{aligned} & \frac{f(x_1)}{(x_1-x_2)(x_1-x_3)\dots(x_1-x_n)} + \frac{f(x_2)}{(x_2-x_1)(x_2-x_3)\dots(x_2-x_n)} + \dots \\ & \qquad \qquad \qquad + \frac{f(x_n)}{(x_n-x_1)(x_n-x_2)\dots(x_n-x_{n-1})}. \end{aligned}$$

$$\text{Let } \frac{f(x)}{(x-x_1)(x-x_2)\dots(x-x_n)} \equiv \frac{A_1}{x-x_1} + \frac{A_2}{x-x_2} + \dots + \frac{A_n}{x-x_n}.$$

Then $A_1, A_2, A_3, \dots, A_n$

$$\begin{aligned} &= \frac{f(x_1)}{(x_1-x_2)(x_1-x_3)\dots(x_1-x_n)}, \quad \frac{f(x_2)}{(x_2-x_1)(x_2-x_3)\dots(x_2-x_n)}, \quad \dots, \\ & \qquad \qquad \qquad \frac{f(x_n)}{(x_n-x_1)(x_n-x_2)\dots(x_n-x_{n-1})}. \end{aligned}$$

Hence, $f(x) \equiv \sum A_1(x-x_2)(x-x_3)\dots(x-x_n)$ = polynomial of degree $\geq n-2$.

Hence, $\sum A_1 = 0$.

This problem, as we thought, proves to be similar to Ex. 4, p. 319, 3rd Edition of Burnside and Panton's *Theory of Equations*. ED. F.

GEOMETRY.

311. Proposed by J. OWEN MAHONEY, B. E., M. Sc., Dallas High School, Dallas, Texas.

Triangle ABC is obtuse-angled at C ; x, y, z are squares on the sides AC, CB, BA ; LH and MJ are lines joining adjacent sides of x, z and y, z . The common chord of the circles on LH and MJ as diameters passes through C and the mid-point of HJ .